ATTORNEY'S DOCKET NUMBER

33146RD004

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING DATE PRIORITY DATE CLAIMED PCT/US99/05733 5773 17 March 1999 18 March 1998

TITLE OF INVENTION APPLICANT(S) FOR DO/EO/US

"PROCESS FOR REDUCING PROTEIN ALLERGENS IN LATEX PRODUCTS"

George W. WEINHERT

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

- This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
- □ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
- This express request to begin national examination procedures (35 U.S.C. 371(f) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(l).
- A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
- A copy of the International Application as filed (35 U.S.C. 371(c)(2))
- a. Is transmitted herewith (required only if not transmitted by the International Bureau).
- b. □ has been transmitted by the International Bureau.
- c. ⋈ is not required, as the application was filed in the United States Receiving Office (RO/US).
- ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
- a, a are transmitted herewith (required only if not transmitted by the International Bureau).
- b. I have been transmitted by the International Bureau.
- c. \square have not been made; however, the time limit for making such amendments has NOT expired.
- d. A have not been made and will not be made.
- 8.

 A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
- 9.

 An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
- 10.

 A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

- 11.

 An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
- 12.

 An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
- 13.

 A FIRST preliminary amendment.
 - A SECOND or SUBSEQUENT preliminary amendment.
- 14.

 A substitute specification.
- ¶5. □ A change of power of attorney and/or address letter.
- a. First page of WO 99/47087 publication
- PCT/ISA/210

2000 27 2007

THE RESERVE AND THE

, and

Harry Street

13



DESIGNATED/ELECTED OFFICE (DO/EO/US)

U.S. DEPARTMENT OF COMMERCE; PATENT AND TRADEMARK

ATTORNEY'S DOCKET NUMBER

33146RD004

U.S. APPLICATION NO. (if known,

co	U.S. APPLICATION NO. (if known, see 1370 ft 15) 646415				
17. The following fee	CALCULATIONS	PTO USE ONLY			
Basic National Fee (Search Report has b					
International prelimi (37 CFR 1.482)					
No international presearch fee paid to U					
Neither international international search		:			
International preliminand all claims satisf					
	E	NTER APPROPRIATE BA	ASIC FEE AMOUNT =	\$96.00	
Surcharge of \$130.00 for the earliest claimed prior	\$				
Claims	Number Filed	Number Extra	Rate		
Total Claims	20-20 =	00	x \$18.00		•
<u>independent</u> Claims	5 -3 =	2	x \$78.00	\$156.00	
Multiple dependent clain	n(s) (if applicable)		+ \$260.00	0.00	
		TOTAL OF ABOV	E CALCULATIONS =	\$ 252.00	
Heduction by ½ for filing	\$0.00				
			SUBTOTAL =	\$ 252.00	
Processing fee of \$130. from the earliest claimed	20 □ 30 months +				
*.	\$252.00				
Fee for recording the end accompanied by an appr	0.00				
*	\$252.00				
				Amount to be refunded	\$
				charged	\$
a. A check in the amount of \$ 252.00 to cover the above fees is enclosed.					
b. □ Please charge my Deposit Account No in the amount of \$ to cover the above fees.					
 c. \[\begin{align*}					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					

SEND ALL CORRESPONDENCE TO:

SMITH, GAMBRELL & RUSSELL, LLP The Beveridge, DeGrandi, Weilacher & Young Intellectual Property Group 1850 M Street, N.W., Suite 800,

Washington, D.C. 20036 Telephone: (202) 659-2811 Facsimile: (202) 659-1462

SIGNATURE

Robert G. Weilacher, Reg. No. 20,531
NAME REGISTRATION NO.

Date: September 18, 2000

PROCESS FOR REDUCING PROTEIN ALLERGENS IN LATEX PRODUCTS

Reference to Related Application

This application claims priority on U.S. 60/078,388 filed on March 18, 1998 which is incorporated herein in its entirety.

Industrial Applicability

This invention relates to a process for reducing protein allergy caused by latex products, such as latex gloves and latex-containing devices used by the medical profession, which are made from the latex sap of the Hevea brasilisensis tree.

Background of the Invention

It is well known that latex products, such as latex gloves, condoms, catheters and automobile tires made from the sap of the Hevea brasilisensis tree cause allergic reactions to some individuals who come into contact with such products. The allergic reaction is caused by the water-soluble proteins present in the sap of the tree and in the product made therewith.

In the late 1980s the United States Occupational Safety and Health Administration (OSHA) published Bloodborne Pathogen Standards requiring increased use of gloves to protect health care workers from exposure to the AIDS and hepatitis B viruses. Latex glove production substantially increased prior to and following the publication of such Standards. In 1991,

CS + C + I T . C I I I I I I I

5

20

25

25

5

1 4 3 4 3 \ ? () 93 8 wo 99.27087 109 (

the United States Food and Drug Administration (FDA) issued a latex alert regarding allergic reactions of patients and medical personnel who had come in contact with latex products. Among the latex products identified as potentially hazardous by the FDA were surgeon's gloves, latex exam gloves, latex condoms, barium enema retention rings and Foley catheters. The latex alert was issued after the number of annual cases regarding allergic reactions resulting from latex products increased from a few to 1,600. Of the three types of latex related diseases - dermatitis, cell mediated allergy and systemic allergy - which manifest themselves through different symptoms, cell mediated allergic response is a true allergic

response, with reaction restricted to the area of contact

between the glove and the skin when the glove is made from the

sap of the Hevea brasilisensis tree. The reaction may include

swelling and blistering and, after washing of the hands upon

removal of the gloves, it takes from about 24 to 48 hours for

the person's skin to return to normal. This allergy is caused

by several water soluble proteins in latex sap.

Systemic latex allergy, the most serious of the latexrelated diseases, is characterized by allergic rhinitis

- asthma - and can escalate to anaphylaxis and death. This
aflergy is caused by several water soluble proteins in the
latex sap from the Hevea brasilisensis tree present in
products made from such sap, which proteins are dispersed in
the air and are breathed in by people. Methods of measurement

25

5

of airborne concentrations of latex antigen as well as latex antigen concentrations in products, such as gloves, in amount as low as 1 nanogram per cubic meter (ng/M^3) are known. Several healthcare institutions have decided to adopt 10 ng/M^3 as a ceiling concentration or standard for personal exposure.

It is known that up to 100 percent of the water soluble protein can be removed from latex of the Hevea brasilisensis tree by subjecting the latex to several washings with water and to centrifugation, but with each washing the latex yield decreases, thus increasing the cost of the resulting purified product. Numerous attempts have been made by others to solve the problem of allergic reactions caused to certain people when they come into contact with rubber products made from the sap of the Hevea brasilisensis tree due to the proteins in the sap being present in such products.

U.S. Patent No. 5,580,942 acknowledges the severe allergic reactions in hypersensitive people caused by the proteins present in the natural rubber latex obtained from Hevea brasilisensis trees. The patentee has a simple solution to the problem, namely, avoiding use of the latex from the Hevea brasilisensis in making latex products. Instead, the patentee uses latex extracted from the Parthenium argontatum (guayule) plant or the Ficus elastica plant, which plants have a different protein profile, whereby the proteins from the sap of these plants do not cause allergic reactions in hypersensitive humans.

25

5

-4-

A review of latex measurement proteins has been published (Beezhold D.H., Measurement of latex proteins by chemical and immunological methods, Procedings of Latex Protein Allergy:

The Present Position. Amsterdam, Dec. 1993).

Proteins have been isolated from rubber particles and from the B and C serum fractions of fresh non-ammoniated latex (NAL). When analyzed by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE), distant protein bands are observed (Hasma, J. Nat. Rubb. Res., 1992, 7(2), 102-112; Arrequin, et al., Electrophoresis, 1988, 9, 323-326; Slater, et al., J. Allergy, Clin. Immunolo., 1992, 89, 673-678.). Electrophoretic profiles of NAL reveals major proteins at 46, 29 and 14kDa, and minor bands at 90, 55, 40, 36, 24, 20 and 18kDa. Hevein (4kDa) and hevamine (29kDa) are two proteins in latex sera which have been cloned and sequenced (Lee et al., Biol Chem., 1991, 256, 15944-15948; Jekel et al., Eur. J. Biochem., 1991, 200, 123-130). In addition, rubber particles contain the tightly bound proteins prenyltransferase (38kDa) and rubber elongation factor (14.5kDa) which have also been sequenced (Dennis et al., Biol. Chem., 1989, 264, 18618-Light et al., J. Biol. Chem., 1989 264, 18589-97).

Ammonification of latex alters the proteins. Storage of latex in ammonia alters the electrophoretic profiles of proteins such that SDS-PAGE profile changes from distinct bands to a smear of polypeptides with an increase in high molecular weight material (Breezhold et al., Arch. Surg.,

25

5

1992, 127, 1354-1357). Many, but not all, of the changes that occur are due to hydrolysis of the proteins. Since NAL serum proteins migrate primarily below 46kDa, the appearance of high molecular weight polypeptides suggests that ammonification (and/or other compounding ingredients) induce a type of polymerization of the latex proteins that produces the larger polypeptides. This process may contribute to the allergenicity of the latex proteins. In addition, ammonification also extracts some of the rubber bound proteins (primarily rubber elongation factor) making them soluble proteins and thus potential allergens (Hasma, J. Nat. Rubb. Res., 1992, 7(2), 102-112).

Much of the protein in latex is not tightly bound to the rubber, but is "water soluble" and readily leaches out of the latex. In order to measure latex proteins it is important to understand the parameters which influence extractability. It has been shown that water soluble proteins are readily extractable, however, complete extraction may take up to 18 hours or more (Dalrymple et al., Rubb. Devel., 1992, 45, 51-60; Hashim, International Rubber Technology Conference, Kuala Lumpur, Malaysia, June 1993; Yeang et al., International Rubber Technology Conference, Kuala Lumpur, Malaysia, June 1993). The volume, pH, and composition of extraction buffer are also important factors.

25

5

A majority of the latex proteins have an acidic pH between 4.0 and 6.5 (Chambeyron et al., Allergy, 1992, 90, 230-235) that increases their solubility in basic buffers and helps explain the observation that more protein is extracted in higher pH buffers. Latex proteins are drawn to the surface of the latex during drying. The surface proteins can be collected directly from the surface by dry swabbing (Dalrymple et al., Rubb. Devel., 1992, 45, 51-60). By analyzing proteins obtained from dry swabbing rubber films, a unique group of surface proteins which are nearly insoluble in water and have limited solubility in carbonate buffer was observed. proteins migrate in SDS gels with a relative molecular mass of between 60 and 70kDa. The proteins are remarkable in that they can be identified by their ability to non-specifically bind IqM from human serum. Furthermore, the IgM binding proteins (IgMbp) can activate the serum complement system and thereby have the potential to cause anaphylactoid reactions. Because these proteins are insoluble they are easily overlooked but must be considered as a potential source of allergens.

Attempts to identify specific allergens has resulted in a wide range of different molecular weight allergens being proposed (Chambeyron, et al., Allergy, 1992, 47, 92-97;

Makinen-Kiljunen et al., J. Allergy Clin. Immunol., 1992, 90, 230-235; Jaeger et al., Allergy Clin. Immunol., 1992, 89,

25

759-768). A recent review summarized the published data and suggested that the studies implicate two proteins of 30kDa and 14kDa as the common allergens (Hamann, Amer. J. Cont. Derma., 1993, 4, 4-21). An abstract recently identified a 14kDa allergen as rubber elongation factor, one of the rubber bound proteins (Czuppon et al., CHEST 104; abstract 159S, 1993).

- U.S. Patent No. 5,610,212 acknowledges that products made of natural rubber, such as rubber gloves, produce allergic reactions in some people, which reactions are attributed to the proteins present in natural rubber. The patent also discusses prior deproteinizing processes for latex which have been used to get around the problem caused by natural rubber. The patent discloses a process for markedly improving the stabilization of deproteinized natural rubber latex which has been treated with a protease and a surfactant, by addition thereto of a specific surfactant or an oligomer or polymer.
- U.S. Patent No. 5,622,998 discusses various known processes for depolymerizating natural rubber and discloses a process for forming a liquid depolymerized natural rubber which produces no immediate allergy. The patentee dissolves a deproteinized natural rubber into an organic solvent to a concentration of about 1 to 30 percent by weight and then carries out air oxidation of the resulting solution in the presence of a metallic catalyst.

25

5

-8-

Brief Summary of the Invention

The process of the invention comprises reacting proteins in the sap of the Hevea brasilisensis plant and other rubber plants, which sap is used in the known processes for the manufacture of latex products, such as latex gloves or other latex-containing products, automobile tires and medical devices which are intended to come into contact with people or will be exposed to people, such as patients, doctors, nurses, laboratory technicians and others, with an aldehyde, such as a mono-aldehyde, such as formaldehyde, or a dialdehyde, such as glutaraldehyde or semialdehydes or any chemical containing an aldehyde group, to cross-link such proteins. The cross-linked proteins no longer have the capability to cause an allergic reaction to persons coming into contact with the latex products made by the process of the invention. The crosslinking reaction between the proteins in the latex sap and the aldehyde can take place in the solution used to prepare the final product, or after the final latex product has been formed, or during various intermediate steps of the known processes for forming the latex products.

Detailed Description of the Invention

The invention comprises a process for significantly reducing and/or eliminating the allergy caused to certain people who come into contact with latex products made from the latex sap of the Hevea brasilisensis plant or any latex

25

5

producing plant due to the presence of proteins from the latex sap. Processes for making such latex products are well known. Applicant has found that he can significantly decrease and/or even eliminate the presence of these proteins in the sap which causes the allergic reactions by reacting such proteins with an aldehyde, such as formaldehyde, or preferably with a dialdehyde, most preferably, glutaraldehyde. Such reaction can take place in the latex solution used in a step of the known process of making the latex product. Such aldehyde, and preferably glutaraldehyde, is added to and stirred in the latex solution in an amount sufficient to react with the proteins in the latex solution and to cross-link substantially all of the proteins into polymers. Alternatively, the reaction between the proteins in the sap of the latex and the aldehyde can take place during other steps in the known processes for making the latex products, as long as the proteins in the latex come into contact with and react with the aldehyde and are cross-linked to form a polymer. Such reaction can also take place after the latex product, such as the rubber glove, is formed. The reaction of the proteins in the latex which are on the surfaces of the latex product are reacted with the aldehyde, such as by the coating of the surface with the aldehyde, or by the immersion of the product in the aldehyde, for a period of time sufficient to completely react and cross-link the proteins with the aldehyde.

25

5

-10-

The use of an aldehyde and, particularly, glutaraldehyde, to obtain the significant reduction of protein allergies caused by latex products made from the sap of the Hevea brasilisensis tree is contrary to what one would be expected to use. Glutaraldehyde is one of two high-level disinfectant chemicals currently approved by the FDA for disinfecting endoscopes, bronchoscopes, cystoscopes, ultrasonic transducers and other devices not amenable to disinfection by heat, steam, radiation or other means. Glutaraldehyde presents a health hazard to persons working with it, such as red burning eyes, sore throat, nasal discharge and red itchy skin. Glutaraldehyde is also a skin sensitizer and has been known to aggravate asthma. Applicant has found that by reacting two known health hazards to humans, namely, the proteins in the latex sap of the Hevea brasilisensis tree and glutaraldehyde, the proteins cross-linked with the glutaraldehyde form a reaction product which eliminates the protein allergy of the latex product.

The latex with reduced allergans that is produced by the method of the invention will result in significantly increased safety for latex allergic persons who use or are exposed to the following types of latex containing products:

Emergency Equipment

Blood pressure cuffs, stethoscopes, disposable gloves, oral and nasal airways, endotracheal tubes,

25

5

-11-

tourniquets, intravenous tubing, syringes and electrode pads

Personal Protective Equipment

Gloves, surgical masks, goggles, respirators and rubber aprons

Office Supplies

Rubber bands and erasers

Hospital Supplies

Anesthesia masks, catheters, wound drains, injection ports, rubber tops of multidose vials and dental dams

Consumer Products

Automobile tires, motorcycle and bicycle handgrips, carpeting, swimming goggles, racquet handles, shoe soles, expandable fabric (waistbands), dishwashing gloves, hot water bottles, condoms, diaphragms, balloons, pacifiers, and baby bottle nipples

The following examples are merely illustrative of the process of the invention.

The radio-immunosorbant assay (RAST) is often used as a sensitive technique which employs radio-isotope labelled anti-IgE to measure specific IgE antibody in patient sera. The latex RAST (Pharmacia) is used to determine if specific IgE to the latex proteins is present and is semi-quantitative in

25

5

-12-

determining the amount of IgE present. The assay is used primarily as a diagnostic test for latex allergy.

The RAST assay employed below is performed as a competition assay (RAST inhibition) to quantitate the amount of allergen in a latex extract. In this assay, soluble allergens in latex extracts compete for binding to latex specific IgE in pooled sera from latex allergic individuals. When soluble allergens react with the IgE, the antibody is prevented from binding to a solid phase latex allergen preparation. The amount of inhibition reflects the level of soluble allergens in the extract. The RAST inhibition assay is a very sensitive method to quantitate latex allergens.

Examples of Latex Allergan Reduction by the Process of the Invention

Example 1.

1. Ammoniated latex (C.N.L from the General Latex and Chemical Corporation, Billerica, MA) was treated with glutaraldehyde and formaldehyde to effectively cross-link the natural rubber proteins. 20 cc of a 20 % glutaraldehyde solution (Sigma Chemical, St Louis, MO, #G6257) were added to 700 cc of latex sap. 22 cc of a 37% solution of formaldehyde (Fisher Chemical, Fair Lawn, NJ) were added to 700 cc of latex sap. Both experiments and a control of latex sap were conducted at room temperature, which was 23 degrees centigrade (C). Ten minutes later, the formaldehyde-latex solution became gel like,

-13-

indicating the cross-linking of protein had occurred. The glutaraldehyde-latex solution did not become gel-like.

- 2. The glutaraldehyde-latex solution was coated on a glass plate and allowed to air dry (23 degrees C for 30 minutes). A control of the untreated latex was applied to another glass plate in a similar fashion and allowed to air dry (23 degrees C for 30 minutes). The plates were incubated at 130 degrees C for 30 minutes.
- 3. The formed films were sent to the MAYO clinic to determine the concentration antigen by the RAST inhibition test, a test specific to measure the antigenic proteins. The control sample contained 409,140 nanograms per gram of antigenic protein, and the glutaraldehyde treated latex sample contained 42,672 nanograms per gram of antigenic protein.
- 4. Subsequent samples were prepared in a similar fashion described above, except the vulcanization temperature was 120 degrees centigrade. The RAST inhibition antigen testing was conducted at the IBT Reference Laboratory. The control sample contained 13.0 micrograms per gram of antigenic protein, and the glutaraldehyde treated latex sample contained 5.6 micrograms per gram of antigenic protein.

20

Example 2.

Additional experiments were conducted using a new sample of latex and several other aldehydes.

5

- 1. Ammoniated latex (C.N.L. from the General Latex Corporation, Billerica, MA) was treated with formaldehyde and glutaraldehyde as confirmation of earlier experimental tests demonstrating the crosslinking of the natural rubber proteins previously discussed. In addition, samples of the latex were treated with, citronellal (Fisher Scientific, Pittsburgh, PA, #AC405291000), butyraldehyde (Fisher Scientific, Pittsburgh, PA, # AC220302500) and crotonaldehyde (Fisher Scientific, Pittsburgh, PA #AC 158220050). An aliquot of each aldehyde was added to 20 milliliters of latex dispersion. All five experiments were conducted at room temperature, which was 22 degrees centigrade (C). There was no apparent thickening in any of the five solutions.
 - 2. Each solution was coated on a glass plate and allowed to air dry for 30 minutes at 22 degrees centigrade (C). The plates were incubated for 20 minutes at 200 degrees Fahrenheit
 - 3. The formed films were sent to the MAYO Clinic to determine the concentration of antigen by the RAST inhibition test, a test specific to measure antigenic proteins. This control sample contained 9370 nanograms per gram of antigenic protein.

10

5

0

5:

- 4. 400 microliters of formaldehyde were added to 20 milliliters of latex. The sample contained 2249 nanograms per gram of antigenic protein. This was a 76 percent reduction as compared to the control sample.
- 5. 800 microliters of glutaraldehyde were added to 20 milliliters of latex, and the sample contained 5254 nanograms per gram of antigenic protein. This was a 44 percent reduction as compared to the control sample.
- 6. 800 microliters of butyraldehyde were added to 20 milliliters of latex, and the sample contained 1461 nanograms per gram of antigenic protein. This was an 84 percent reduction as compared to the control sample.
- 137. 400 microliters of crotonaldehyde were added to 20 milliliters of latex, and the sample contained 5554 nanograms per gram of antigenic protein. This was a 41 percent reduction as compared to the control sample.
 - 8. 200 microliters of citronellal were added to 20 milliliters of latex, and the sample contained 4515 nanograms per gram of antigenic protein. This was a 52 percent reduction as compared to the control sample.

20

15

Example 3.

Examples of Latex Allergan Reduction in a Typical Manufacturing Process

A typical manufacturing sequence for a dipped product such as a glove or a condom involves sequential steps of:

THE STATE OF

- 1. dipping a form or mandrill into a release agent
- 2. air drying the release agent coated form or mandrill
- 3. dipping the form or mandrill into a latex solution
- 4. air drying the latex coated form or mandrill
- 5. vulcanizing the latex, and generally,
- 6. dipping the vulcanized product a dispersion of powder
 The product is then stripped from the form or mandrill and packaged.
 Some products, such as surgical gloves, may also be sterilized,
 usually by radiation.

The method of the invention can be performed at different steps of the manufacturing process. The aldehyde solution can be mixed with the latex dipping solution. Generally the aldehyde is added to the latex solution with mixing to promote uniformity. The aldehyde can also be in line blended with the latex solution to maintain a consistent depth of the dipping solution. Currently, some glove manufactures spray water on the vulcanized glove to wash away water-soluble surface protein or pass the gloves on the mandrills through a series of leaching tanks. By the method of the invention, the aldehyde can be added to the washing or leaching solutions or could

-17-

be sprayed directly onto the finished product to cross-link the surface proteins.

The above examples demonstrate that substantial reductions in latex antigens can be obtained by the method of the invention.

Further variations and modifications will be apparent to those skilled in the art and are intended to be encompassed by the claims appended hereto.

graph of the same

I CLAIM:

 A process for reducing the antigenicity of proteins in latex sap comprising:

contacting a sufficient amount of an aldehyde with an antigenic latex protein to react with and cross-link the protein so as to significantly reduce the antigenicity of the protein.

2. A process for reducing the antigenicity of proteins in latex sap comprising:

forming a solution of a latex sap containing an antigenic protein, and

adding a sufficient amount of an aldehyde to react with and cross-link the protein so as to significantly reduce the antigenicity of the protein.

- 3. A latex sap produced by the process according to claim 2.
- 4. The process as defined in Claim 1, wherein said aldehyde is a mono-aldehyde or a dialdehyde.
- 5. The process as defined in Claim 3, wherein said dialdehyde is glutaraldehyde.
- 6. The process as defined in Claim 3, wherein said monoaldehyde is formaldehyde.

7. A process for making a latex product with reduced antigenicity from latex sap containing antigenic proteins, comprising:

forming a solution of said latex sap,

applying a sufficient amount of an aldehyde to a

surface of the product to react with and cross-link the protein so
as to significantly reduce the antigenicity of the protein,

shaping said latex sap into a product, and

vulcanizing said product.

- 8. The process as defined in Claim 7, wherein said aldehyde is a mono-aldehyde or a dialdehyde.
- 9. The process as defined in Claim 8, wherein said dialdehyde is glutaraldehyde.
- 10. The process as defined in Claim 8 wherein said monoaldehyde is formaldehyde.
- 11. A process for making a latex product with reduced antigenicity from latex sap containing antigenic proteins, comprising:

forming a solution of said latex sap, shaping said latex sap into a product, vulcanizing said product, and

applying a sufficient amount of an aldehyde to a surface of the product to react with and cross-link the protein so as to significantly reduce the antigenicity of the protein.

- 12. The process as defined in Claim 11, wherein said aldehyde is a mono-aldehyde or a dialdehyde.
- 13. The process as defined in Claim 12, wherein said dialdehyde is glutaraldehyde.
- 14. The process as defined in Claim 12 wherein said monoaldehyde is formaldehyde.
- 15. A latex product produced by the process according to claim7.
- 16. A latex product produced by the process according to claim11.
- 17. In a process for making a latex product from the latex sap of the Hevea brasilisensis plant containing antigenic proteins which cause allergic reactions to persons coming into contact with said latex product, comprising;

forming a solution of said latex sap, shaping said latex sap into a product and vulcanizing said product, the improvement comprising the step of reacting said proteins

present in said latex sap with a sufficient amount of an aldehyde to react with and cross-link said proteins, so as to significantly reduce the allergenicity of said latex product.

- 18. The process as defined in Claim 17, wherein said aldehyde is a mono-aldehyde or a dialdehyde.
- 19. The process as defined in Claim 18, wherein said dialdehyde is glutaraldehyde.
- 20. The process as defined in Claim 18, wherein said monoaldehyde is formaldehyde.

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISI	HED U	JNDER THE PATENT COOPERATION TREATY (PCT)
(51) International Patent Classification ⁶ :		(11) International Publication Number: WO 99/47087
A61F 6/00, A61L 2/00, 27/00, 31/00, C08F 34/00, 134/00	A1	(43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/USS (22) International Filing Date: 17 March 1999 (1800) (30) Priority Data: 60/078,388 18 March 1998 (1803.98) (71) Applicant (for all designated States except US): RISS AGEMENT ASSOCIATES LTD. [US/US]; 62 Pois Worcester, MA 01604 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): WEINERT, Georgius [US/US]; 62 Pointe Rok, Worcester, MA 01604 (US). (74) Agents: DEGRANDI, Joseph, A. et al.; Smith, Georgius Russell, LLP, The Beveridge, DeGrandi, Weil Young Intellectual Property Group, Suite 800, Street, N.W., Washington, DC 20036 (US).	UK MANinte Roorge, V JS). Gambre lacher 1	BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, IP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: PROCESS FOR REDUCING PROTEIN ALLE (57) Abstract	EKGEN	S IN LATEX PRODUCTS

Described is a process for reducing the antigenicity of sap and products made from the sap of the *Hevea brasilisensis* plant and other rubber plants. The process involves contacting sap or a latex rubber product with a mono or dialdehyde, a semialdehyde or any chemical containing an aldehyde group, to cross—link antigenic proteins within the sap or the latex product. The cross—linked proteins no longer have the capability to cause an allergic reaction to persons coming into contact with the latex products made by the process of the invention. The cross-linking reaction between the proteins in the latex sap and the aldehyde can take place in the solution used to prepare the final product, or after the final latex product has been formed, or during various intermediate steps of the processes for forming the latex products.

Declaration and Power of Attorney United States Patent Application

Patents and Design Patents
Sole & Joint Inventors
Sole & Non-convention
PCT & Non-PCT
This form cannot be amended, altered
or changed after it is signed
(For use only for inventors who
understand the English language.)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on

the invention entitled Process For Reducing Protein Allergens In Latex Products

(check one)	□ is attached he⋈ was filed as U.	ereto. S. Application No. 09/646,415	on <u>September 18, 2000</u> and (if a	pplicable) was amended on
	was filed as I	PCT International Application No ler PCT Article 34 on	()	ch 17, 1999 and (if applicable) was
I have reviewed I acknowledge t	I and understand the cont the duty to disclose infor	ents of the above-identified specification, in mation which is material to patentability as	ncluding the claims, as amended by any ame defined in Title 37, Code of Federal Regula	endment referred to above. ations, §1.56.
have also identi	oob(a) of any PC1 intern	ational application which designated at lear oplication for patent or inventor's certificate	e-(d) or §365(b) of any foreign and PCT app st one country other than the United States of or PCT international application having a	of Amounton lines of its ship Double at the
Foreign/l	PCT Application No.	Country	Filing Date	Priority Claimed? (yes/no)
application or Point in the state of application which application and	America listed in this De CT international applicate ch is material to patental	claration and, insofar as the subject matter ion in the manner provided by the first para	any United States application and PCT intent of each of the claims of this application is r agraph of Title 35, United States Code, §11 I Regulations, §1.56 which became availab	not disclosed in the prior United States
CHÉCHE:	Application No.	Filing Date	Status (patented	/pending/abandoned?)
TOTAL STATE OF THE				
I horaby alaim n	ministration of the state of th	05TI 1 101 - 0 1 0100 - 0 -		
U.S. Provis	sional Application No.	e 35 United States Code §119(e) of any U. Filing Date March 18, 1998	S. provisional application(s) listed below:	·
(32,263), Dennis	31), Herbert M. Hanegar	i (25,682), Dale Lischer (28,438), Frederic Villiam F. Rauchholz (34,701), Michael C.	all business in the Patent and Trademark Ock F. Calvetti (28,557), J. Rodgers Lunsford Carrier (42,391), Eric J. Hanson (44,738),	1 III (29 405) Michael A Makuch
Send all correspondence 263-4329. Direct	ondence to: SMITH, GAI ct all phone calls to (202)	MBRELL & RUSSELL, LLP, 1850 M Stre 0 659-2811.	eet, N.W. (Suite 800), Washington, D.C. 20	036. All facsimiles may be sent to (202)
both, under Secti	ier that these statements i	vere made with the knowledge that willful e United States Code and that such willful	hat all statements made on information and false statements and the like so made are pu false statements may jeopardize the validity	mishable by fine or imprisonment or
Residence (ci	ity, state, country):	or: George W. WEINERT Pocasset, MA VVVV int Drive, Pocasset, MA 02559 1	Citizenship: USA	
Signature:	George W	Win	Date: ZI MARCH 2	002
	second joint invent ty, state, country): ldress:	or, if any:		Citizenship:
Signature:	- N		Date:	
- A 11'4' 1'				

United States Patent & Trademark Office Office of Initial Patent Examination -- Scanning Division



Page(s) for scanning.	of (Document title)	were not present
☐ Page(s)for scanning.	of(Document title)	were not present

□ Scanned copy is best available.